

THE OVERHAUSER INSTABILITY

Zoltán Radnai and Richard Needs
TCM Group

ESDG Talk
14th February 2007


INTRODUCTION

- Hartree-Fock theory and Homogeneous Electron Gas
- Noncollinear spins and Spiral Spin Density Waves
- The Overhauser Instability
- Results
- Interpretation

NONCOLLINEAR SPINS

- *Collinear*: definite spin (up or down) with respect to global quantization axis.
- Particles are distinguishable, simplifies calculation.
- *Noncollinear*: spin directions that are not parallel to the global quantization axis and the spin direction can vary with position.
- Wavefunction depends on position and spin coordinates.

Spin density or Magnetization density: net magnetic moment due to spin.
Vector function of position. Noncollinear example:



Spiral Spin Density Wave
(magnetization wavevector \mathbf{q})

GENERALIZED HARTREE-FOCK THEORY

- HF theory of noncollinear spins. Orbitals have full spin dependence.
- Can write orbitals as function of space-spin coordinates $\psi(\mathbf{x})$.
- Can write orbitals as two-component spinors: $\underline{\psi}(\mathbf{r}) = \begin{pmatrix} \psi_1(\mathbf{r}) \\ \psi_2(\mathbf{r}) \end{pmatrix}$.
- Determinant of spinors: $\Psi = \frac{1}{\sqrt{N!}} \det |\underline{\psi}_i(\mathbf{r}_j)|$.
- Total energy is $E = \langle \Psi | \hat{H} | \Psi \rangle$. Evaluate variation with respect to orbitals, subject to orthonormality constraint.

- Gives single-particle HF equation (depending on form of \hat{H}) such as:

$$(\hat{\mathbf{K}} + \hat{\mathbf{U}} + \hat{\mathbf{V}} - \hat{\mathbf{J}})\underline{\psi}_k(\mathbf{r}) = \epsilon_k \underline{\psi}_k(\mathbf{r}) \quad (1)$$

- This is a 2x2 matrix equation. $\hat{\mathbf{K}}$ (kinetic energy), $\hat{\mathbf{U}}$ (external potential), $\hat{\mathbf{V}}$ (direct term) and $\hat{\mathbf{J}}$ (exchange term) are 2x2 matrices of spatial operators.
- For spin independent \hat{H} , $\hat{\mathbf{J}}$ can still have off-diagonal components, giving rise to noncollinearity.
- Collinear case is special case with $\underline{\psi}(\mathbf{r}) = \phi(\mathbf{r})\underline{\chi}$, where $\underline{\chi}$ is a spin eigenstate. Reduces problem to Unrestricted HF theory.
- HF equation needs to be solved self-consistently.

OVERHAUSER INSTABILITY

- What is ground state of Homogeneous Electron Gas in Hartree-Fock theory?
- High-density (low r_s) limit: paramagnet. Low-density (high r_s): ferromagnet. Collinear states.
- Overhauser: paramagnet is never ground state, but instead a Spiral Spin Density Wave is.
- Proof and analytical solution for 1D with repulsive δ -function interaction. (A. W. Overhauser, *Giant spin density waves*, Phys. Rev. Lett. **4**, p.462 (1960))
- Proof of existence for 3D with Coulomb interaction. (A. W. Overhauser, *Spin density waves in an electron gas*, Phys. Rev. **128**, p.1437 (1962))

HARTREE-FOCK THEORY OF HOMOGENEOUS ELECTRON GAS


- Hamiltonian: $\hat{H} = \sum_i -\frac{1}{2}\nabla_i^2 + \sum_i U(\mathbf{r}_i) + \frac{1}{2} \sum_{i,j \neq i} V(\mathbf{r}_i, \mathbf{r}_j) + \lambda$
 V could be Coulomb or Ewald interaction.

- Single-particle HF equation is self-consistently solved by SSDW orbitals:

$$\underline{\psi}_{\mathbf{k}}(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}} \begin{pmatrix} \cos(\frac{1}{2}\theta_{\mathbf{k}}) e^{-i\frac{1}{2}\mathbf{q}\cdot\mathbf{r}} \\ \sin(\frac{1}{2}\theta_{\mathbf{k}}) e^{+i\frac{1}{2}\mathbf{q}\cdot\mathbf{r}} \end{pmatrix}$$

- \mathbf{k} is plane-wave vector, \mathbf{q} is magnetization wave vector (constant), and the orbital has its spin pointing in $(\theta, \mathbf{q} \cdot \mathbf{r})$ direction (spiral in space).
- *Mixing* up spin at $\mathbf{k} - \frac{1}{2}\mathbf{q}$ with down spin at $\mathbf{k} + \frac{1}{2}\mathbf{q}$.

- Paramagnet and ferromagnet are special cases. They correspond to particular choices of occupation and $\theta_{\mathbf{k}}$.
- Example: paramagnet is two overlapping spheres in \mathbf{k} -space:



ANALYTICAL PART

- Total energy:

$$E = \sum_{\mathbf{k}} \frac{1}{2} \left\{ \mathbf{k}^2 + \frac{1}{4} \mathbf{q}^2 - \mathbf{k} \cdot \mathbf{q} \cos \theta_{\mathbf{k}} \right\} - \frac{1}{2} \frac{1}{\Omega} \sum_{\mathbf{k}, \mathbf{k}' \neq \mathbf{k}} \frac{4\pi}{|\mathbf{k} - \mathbf{k}'|^2} \cos^2 \frac{1}{2} (\theta_{\mathbf{k}} - \theta_{\mathbf{k}'}) + \frac{1}{2} N \xi \quad (2)$$

(ξ is Ewald self-image term)


- Single-particle HF equation:

$$\left[\begin{pmatrix} K_{\mathbf{k}1} & 0 \\ 0 & K_{\mathbf{k}2} \end{pmatrix} - \begin{pmatrix} J_{\mathbf{k}1} & J_{\mathbf{k}0} \\ J_{\mathbf{k}0} & J_{\mathbf{k}2} \end{pmatrix} \right] \begin{pmatrix} \cos(\frac{1}{2}\theta_{\mathbf{k}}) \\ \sin(\frac{1}{2}\theta_{\mathbf{k}}) \end{pmatrix} = \epsilon_{\mathbf{k}} \begin{pmatrix} \cos(\frac{1}{2}\theta_{\mathbf{k}}) \\ \sin(\frac{1}{2}\theta_{\mathbf{k}}) \end{pmatrix} \quad (3)$$

NUMERICAL PART


- **Aim** Find occupation of \mathbf{k} -space and form of $\theta_{\mathbf{k}}$ self-consistently. Find value of q that gives lowest energy.
- Start by choosing initial occupation of orbitals and $\theta_{\mathbf{k}}$.
- Two schemes: best guess at solution (Overhauser) and paramagnetic occupation with randomized angles.
- Iterate to self-consistency.
- Consistency of result: total energy vs. sum of eigenvalues.
- Numerically tricky, false local minima.

RESULTS: ENERGY VS. SYSTEM SIZE



RESULTS: ENERGY VS. SYSTEM SIZE


SDW HF Extrapolation



Extrapolating to infinite system size


RESULTS: ENERGY VS. Q

SDW HF
r_s 1, N 35114




Lowest energy is achieved when q is near $2k_F$ (3.84 a.u.).

RESULTS: ENERGY VS. r_s




Effect gets smaller at higher density, tricky at $r_s = 1.0$.

NATURE OF THE INSTABILITY



Occupation in \mathbf{k} -space at start




Occupation in \mathbf{k} -space at end

NATURE OF THE INSTABILITY


SDW HF Angles

N10082 rs 5 q18 ky=0 kz=0



NATURE OF THE INSTABILITY

Instability is associated with formation of a gap. The spiral spin density is a short-wavelength periodic structure.



CONCLUSIONS

- Demonstrated the existence of the instability for a range of densities. Its nature is qualitatively as predicted by Overhauser.
- There is always a Spiral SDW solution that has lower energy than paramagnet.
- Work in progress: other types of instabilities, densities below the para/ferromagnetic crossover.
- Work in progress: Quantum Monte Carlo (VMC) calculations using the HF orbitals.

ACKNOWLEDGMENTS

- Financial support was provided by the Gates Cambridge Trust and Trinity College.
- Computational resources were provided by TCM Group.